

Application/Control Number: 10/568,004  
Art Unit: 4153  
July 24, 2008  
Page 2

RECEIVED  
CENTRAL FAX CENTER

JUL 24 2008

**Amendments to the claims**

1-10 (Canceled)

11. (Currently amended) A device for detecting arterial pressure with high measurement precision, comprising a cuff with inflatable chamber, adapted to be placed around the arm of a patient, means for introducing air to inflate said cuff, and decompression means adapted to decompress said inflatable chamber, further comprising means adapted to detect and store all the sphygmic pulses generated by the arterial pulsation and to identify the pulses that correspond to appearance and disappearance of wrist beat, detected by means of a technique for detecting sphygmic pulses generated by arterial pressure that provides for the intervention of an operator to detect the sphygmic pulses and of the operator for a subsequent [[and]] subjective judgment [[of an operator]] of said sphygmic pulses.

12. (Previously presented) The device of claim 11, wherein said decompression means of said inflatable chamber comprise a valve for providing constant and time-controlled decompression.

13. (Previously presented) The device of claim 11, comprising discharge means adapted to completely and instantaneously discharge the inflatable chamber of said cuff.

14. (Previously presented) The device of claim 11, wherein said means for detecting and storing the sphygmic pulses are connected to data storage means, which are adapted to store the chart of the sphygmic pulses.

15. (Previously presented) The device of claim 11, comprising a display that is adapted to display detected levels of pressure and levels of sphygmic intensity of the pulsations.

16. (Previously presented) The device of claim 11, comprising a button that is adapted to be pressed by the operator when the operator detects sphygmic pulses that correspond to systolic or diastolic pressure.

Application/Control Number: 10/568,004  
Art Unit: 4153  
July 24, 2008  
Page 3

17. (Currently amended) A method for detecting arterial pressure, comprising the steps of:

pumping air into a cuff provided with an inflatable chamber;

decompressing said inflatable chamber;

detecting, by means of the intervention and subjective judgment of an operator, using a stethoscope, the sphygmic pulses that correspond respectively to the appearance and disappearance of the wrist beat,

further comprising the steps of:

detecting and storing a chart of all the sphygmic pulses generated by arterial pulsation by using an electronic sensing and storage circuit;

identifying, among said sphygmic pulses, the ones that correspond to the appearance and disappearance of the pulse beat, detected by means of said stethoscope.

18. (Previously presented) The method of claim 17, wherein said step of performing the decompression of said inflatable chamber comprises performing decompression at a controlled and constant rate.

19. (Previously presented) The method of claim 17, further comprising a step of storing said sphygmic pulses generated by arterial pulsation, in order to allow subsequent analysis of the chart of sphygmic pulses, in order to determine assuredly the pulses that actually correspond to the maximum and minimum values of arterial pressure.

20. (Previously presented) The method of claim 17, comprising a step of pressing, on the part of said operator, a button when sphygmic pulses that correspond to systolic and diastolic pressure are detected, said sphygmic pulses that correspond to systolic and diastolic pressure being "marked" on a digital scale of said device.